	Student's Name:	D.O.R.	Teacher	
Place Child				
Picture Her	Asthmatic Yes* No	*Higher risk for severe		
	◆ STEP 1: TRE	ATMENT A		
Symptoms:	<u> </u>	JAINEN V	Give Checke	d Medication**
If a food aller	gen has been ingested, but no symptoms:		☐ Epinephrine	☐ Antihistamine
Mouth	tching, tingling, or swelling of lips, tongo	ie, mouth	☐ Epinephrine	☐ Antihistamine
Skin	Hives, itchy rash, swelling of the face or e	extremities	☐ Epinephrine	☐ Antihistamine
Gut	Nausea, abdominal cramps, vomiting, dia	rrhea	☐ Epinephrine	☐ Antihistamine
Throat†	Fightening of throat, hoarseness, hacking	cough	☐ Epinephrine	☐ Antihistamine
Lung†	Shortness of breath, repetitive coughing, v	wheezing	☐ Epinephrine	☐ Antihistamine
Heart†	Thready pulse, low blood pressure, fainting	g, pale, blueness	☐ Epinephrine	☐ Antihistamine
Other†			☐ Epinephrine	☐ Antihistamine
If reaction is r	rogressing (several of the above areas aff	ected) give:	☐ Epinephrine	☐ Antihistamine
The severity of symp DOSAGE Spinephrine: in see reverse side	toms can quickly change. †Potentially life-threet intramuscularly (circle one) EpiPefor instructions)	eatening.		vinject [™] 0.15 mş
DOSAGE Dinephrine: in see reverse side antihistamine: g	toms can quickly change. †Potentially life-threet intramuscularly (circle one) EpiPe for instructions)	eatening.		vinject [™] 0.15 mş
DOSAGE Dinephrine: in see reverse side antihistamine: g	toms can quickly change. †Potentially life-thr ect intramuscularly (circle one) EpiPe for instructions) tive	eatening. en® EpiPen® Jr. Twir		vinject TM 0.15 mş
DOSAGE Dinephrine: in see reverse side antihistamine: g	toms can quickly change. †Potentially life-thr ect intramuscularly (circle one) EpiPe for instructions) tive	eatening. en® EpiPen® Jr. Twin		vinject™ 0.15 mş
DOSAGE Cpinephrine: in see reverse side Antihistamine: gother: give	toms can quickly change. †Potentially life-thr ect intramuscularly (circle one) EpiPe for instructions) tive	eatening. en® EpiPen® Jr. Twin ion/dose/route	nject™ 0.3 mg Tw	
DOSAGE Cpinephrine: ingsee reverse side Antihistamine: gother: give Call 911 (or Readditional epine	toms can quickly change. †Potentially life-threect intramuscularly (circle one) EpiPerfor instructions) tive	eatening. en® EpiPen® Jr. Twin ion/dose/route GENCY CALLS	aject™ 0.3 mg Tw	
DOSAGE Cpinephrine: ingsee reverse side Intihistamine: g Other: give Call 911 (or Readditional epine	toms can quickly change. †Potentially life-thr ect intramuscularly (circle one) EpiPe for instructions) give	eatening. en® EpiPen® Jr. Twin ion/dose/route GENCY CALLS	aject™ 0.3 mg Tw	
DOSAGE Cpinephrine: ingsee reverse side Intihistamine: gother: give Call 911 (or Readditional epine Dr	toms can quickly change. †Potentially life-thr ect intramuscularly (circle one) EpiPe for instructions) give	eatening. en® EpiPen® Jr. Twin ion/dose/route GENCY CALLS	aject™ 0.3 mg Tw	
DOSAGE Cpinephrine: ingsee reverse side Intihistamine: g Other: give Call 911 (or Readditional epine	toms can quickly change. †Potentially life-thr ect intramuscularly (circle one) EpiPe for instructions) give	eatening. en® EpiPen® Jr. Twin ion/dose/route GENCY CALLS	aject™ 0.3 mg Tw	
DOSAGE Cpinephrine: ingsee reverse side Intihistamine: gother: give Call 911 (or Readditional epine Dr Emergency coname/Relationship	toms can quickly change. †Potentially life-thr ect intramuscularly (circle one) EpiPe for instructions) tive	eatening. en® EpiPen® Jr. Twin ion/dose/route ion/dose/route RGENCY CALLS). State that an allergic Phone Number(s)	nject™ 0.3 mg Tw reaction has been tre 2.)	eated, and
DOSAGE Cpinephrine: ingsee reverse side Intihistamine: gother: give Call 911 (or Readditional epine Dr Emergency coname/Relationship	toms can quickly change. †Potentially life-threect intramuscularly (circle one) EpiPefor instructions) give	eatening. en® EpiPen® Jr. Twin ion/dose/route ion/dose/route RGENCY CALLS). State that an allergic Phone Number(s)	nject™ 0.3 mg Tw reaction has been tre 2.)	eated, and
DOSAGE Cpinephrine: injusee reverse side Antihistamine: g Other: give Call 911 (or Readditional epine Dr Emergency contame/Relationship	toms can quickly change. †Potentially life-thr ect intramuscularly (circle one) EpiPe for instructions) tive	eatening. en® EpiPen® Jr. Twin ion/dose/route RGENCY CALLS). State that an allergic Phone Number(s)	reaction has been tre	eated, and

(Required)

Date_

Doctor's Signature_